

STEDMAN'S Medical Dictionary

26th Edition

ILLUSTRATED IN **COLOR**

Best Available Copy



Williams & Wilkins

Baltimore • Philadelphia • Hong Kong
London • Munich • Sydney • Tokyo

A WAVERLY COMPANY

EX B



Best Available Copy

Editor: Marjory Spraycur
 Senior Editor: Elizabeth Randolph
 Editorial Assistant: Maureen Barlow Pugh
 Copy Editors: Christopher Muldor, Jane Sellman, Barbara Werner
 On-Line Editors: Kathryn J. Cadle, Barbara L. Ferretti, Catherine N. Kelly, Leslie Simpson
 Editorial Proofreaders: Peter W. Binns, Jolanta Obrebska, Carol Sorgen
 Medical Proofreaders: Alfred Jay Bollet, M.D.; John H. Dirckx, M.D.; Thomas W. Filardo, M.D.; Robert Hogan, M.D.; Edward Strim, M.D.
 Database Programmers: Dennis P. Smithers, Dave Marcus, Lexi-Comp Inc., Hudson, OH
 Production Coordinator: Paula K. Huber
 Printing Coordinator: Brian Smith
 Illustration Planning: Wayne J. Hubbel
 Design: Robert C. Och, Dan Pfisterer
 Cover Design: Sharon Reuter, Reuter & Associates

Copyright © 1995
 Williams & Wilkins
 351 W. Camden Street
 Baltimore, MD 21201, USA

Copyright © by William Wood and Company: 1911, 1st ed.; 1912, 2nd ed.; 1914, 3rd ed.; 1916, 4th ed.; 1918, 5th ed.; 1920, 6th ed.; 1922, 7th ed.; 1924, 8th ed.; 1926, 9th ed.; 1928, 10th ed.; 1930, 11th ed.

Copyright © by Williams & Wilkins: 1933, 12th ed.; 1935, 13th ed.; 1939, 14th ed.; 1942, 15th ed.; 1946, 16th ed.; 1949, 17th ed.; 1953, 18th ed.; 1957, 19th ed.; 1961, 20th ed.; 1966, 21st ed.; 1972, 22nd ed.; 1976, 23rd ed.; 1982, 24th ed.; 1990, 25th ed.



All rights reserved. This book is protected by copyright. No part of this book may be reproduced in any form or by any means including photocopying, or utilized by any information storage and retrieval system without written permission from the copyright owner.

Stedman's is a registered trademark of Williams & Wilkins.

Indications, adverse reactions and dosage schedules for drugs set forth in this dictionary are provided by the authors. Williams & Wilkins has independently verified the accuracy of that information and does not make any representation in regard to its accuracy. The reader should review the package information data of the manufacturers of the medications mentioned.

Database design by Lexi-Comp Inc., Hudson, OH
 Printed in the United States of America by R.R. Donnelley & Sons Company

English Language Co-editions
 Asian 1967, 1972, 1976
 Indian 1967, 1973
 Taiwan 1972, 1978

Translated Editions
 Greek 1976
 Indian 1977
 Japanese 1977, 1985, 1995

Portuguese 1976, 1995
 Spanish 1993

Library of Congress Cataloging-in-Publication Data

Stedman, Thomas Lathrop, 1853-1938.

[Medical dictionary]

Stedman's medical dictionary.—26th ed.

p. cm.

ISBN 0-683-07922-0 REGULAR EDITION

ISBN 0-683-07935-2 DELUXE EDITION

1. Medicine—Dictionaries. I. Title. II. Title: Medical dictionary.

[DNLM: 1. Dictionaries, Medical. W 13 S812m 1995]

R121.S8 1995

610'.3—dc20

DNLM/DLC

for Library of Congress

ameloblastic f., a rapidly growing, painful, destructive, radiolucent odontogenic tumor that usually arises through malignant change in the mesenchymal component of a pre-existing ameloblastic fibroma. *syn* ameloblastic sarcoma.

Earle f. f., a transplantable f. derived from subcutaneous tissue of a mouse of C3H strain, grown in tissue culture to which 20-methylcholanthrene had been added.

infantile f., a rapidly growing but infrequently metastasizing f. which usually appears on the extremities in the first year of life.

fi-bröse (fi-brōs'). To form fibrous tissue.

fi-bro-se-rous (fi-brō-sē'ris). Composed of fibrous tissue with a serous surface; denoting any serous membrane.

fi-bro-sis (fi-brō'sis). Formation of fibrous tissue as a reparative or reactive process, as opposed to formation of fibrous tissue as a normal constituent of an organ or tissue.

African endomyocardial f., f. of the inner layers of the myocardium, often including the endocardium, causing diastolic restriction of the heart; indigenous to East Africa.

congenital f. of the extraocular muscles (MIM*135700), an autosomal dominant disorder associated with blepharoptosis and absence of eye movements.

cystic f., cystic f. of the pancreas (MIM*219700), a congenital metabolic disorder, inherited as an autosomal trait, in which secretions of exocrine glands are abnormal; excessively viscous mucus causes obstruction of passageways (including pancreatic and bile ducts, intestines, and bronchi), and the sodium and chloride content of sweat are increased throughout the patient's life; symptoms usually appear in childhood and include meconium ileus, poor growth despite good appetite, malabsorption and foul bulky stools, chronic bronchitis with cough, recurrent pneumonia, bronchiectasis, emphysema, clubbing of the fingers, and salt depletion in hot weather. Detailed genetic mapping and molecular biology have been accomplished by the methods of reverse genetics. *syn* Clarke-Hudfield syndrome, fibrocystic disease of the pancreas, mucoviscidosis, viscidosis.

endocardial f., scarring or collagenosis of the endocardium. *syn* endocardial sclerosis (2).

endomyocardial f., thickening of the ventricular endocardium by f., involving the subendocardial myocardium, and sometimes the atrioventricular valves, with mural thrombosis, leading to progressive right and left ventricular failure with mitral and tricuspid insufficiency; occurs in adults and is endemic in parts of Africa. *syn* Davies' disease, endocardial fibroelastosis (2), endomyocardial fibroelastosis.

idiopathic interstitial f., *syn* usual interstitial pneumonia of Liebow.

idiopathic pulmonary f. (IPF), subacute form also called Hamman-Rich syndrome; an acute to chronic inflammatory process of the lungs, the healing stage of diffuse alveolar damage or acute interstitial pneumonia, either completely idiopathic or associated with collagen-vascular diseases. *syn* chronic fibrosing alveolitis, interstitial pulmonary f.

interstitial pulmonary f., *syn* idiopathic pulmonary f.

leptomeningeal f., a fibrous reaction within the subarachnoid space; sometimes a sequel to infectious or chemical meningitis. *see also* adhesive arachnoiditis.

mediastinal f., f. that may obstruct the superior vena cava, pulmonary arteries, veins, or bronchi; most common cause is histoplasmosis; less commonly tuberculosis or unknown. *syn* fibrosing mediastinitis, idiopathic fibrous mediastinitis.

nodular subepidermal f., *see* dermatofibroma.

oral submucous f., a precancerous condition of the oral mucosa and upper aerodigestive tract characteristically in a native of India.

pericentral f., f. occurring around the central veins in the hepatic lobules.

perimuscular f., f. in the outer media of arteries, usually the renal arteries of young women, where it causes segmental stenosis and hypertension; a variety of fibromuscular dysplasia. *syn* subadventitial f.

pipestem f., a characteristic pipe-shaped f. formed around hepatic portal veins in some cases of long-continued heavy infection with *Schistosoma mansoni*; thought to be induced by the pres-

ence of large numbers of schistosome eggs in the hepatic tissues. *syn* Symmers' clay pipestem f., Symmers' f.

replacement f., the formation of fibrous tissue that occupies sites where various other cells and tissues have become atrophied, or degenerated and necrotic.

retroperitoneal f., f. of retroperitoneal structures commonly involving and often obstructing the ureters; the cause is usually unknown. *syn* idiopathic fibrous retroperitonitis, Ormond's disease, periureteritis plastica.

subadventitial f., *syn* perimuscular f.

Symmers' clay pipestem f., Symmers' f., syn pipestem f.

fi-bro-si-tis (fi-brō-sī'tis). 1. Inflammation of fibrous tissue. 2. Term used to denote aching, soreness, or stiffness, with multiple tender foci (trigger points); unknown etiology; thought by some to be due to a sleep disturbance preventing normal muscle relaxation. *syn* muscular rheumatism. [fibro- + G. -itis, inflammation]

cervical f., *syn* posttraumatic neck syndrome.

fi-bro-tho-rax (fi-brō-thō'raks). Fibrosis of the pleural space.

fi-brot-ic (fi-brō'tik). Pertaining to or characterized by fibrosis.

fi-brous (fi-brōs). Composed of or containing fibroblasts, and also the fibrils and fibers of connective tissue formed by such cells.

fi-bro-xan-tho-ma (fi-brō-zan-thō'mā). A fibrohistiocytic neoplasm.

atypical f., a solitary, often ulcerated, small cutaneous benign tumor composed of foamy histiocytes, spindle cells, and bizarre giant cells; usually found on the exposed skin of older people; microscopically, atypical f. closely resembles malignant fibrous histiocytoma, but originates in the dermis.

fib-u-la (fīb'yū-lā) (NA). The lateral and smaller of the two bones of the leg; it is non-weight bearing and articulates with the tibia above and the tibia and talus below. *syn* calf bone, calf bone (1), perone, peroneal bone, splint bone (2). [L. *fibula* (contr. fr. *figibula*), that which fastens, a clasp, buckle, fr. *figo*, to fix, fasten]

fib-u-lar (fīb'yū-lār). Relating to the fibula. *syn* fibularis (NA). [L. *fibularis*]

fib-u-la-ria (fīb'yū-lā'ria) (NA). *syn* fibular, fibular. [Mod. L.]

fib-u-lo-cal-ca-ne-al (fīb'yū-lō-kāl-kā'nē-āl). Relating to the fibula and the calcaneus.

fi-cin (fī'sin). A proteolytic enzyme isolated from figs (*Ficus carica*, *globata*, and *deliaria*); used in industry as a protein digestant; f. has a wide specificity for protein substrates; an anthelmintic.

Fick, Adolf, German physician, 1829-1901. *see* F. principle, method.

fi-co-sis (fī-kō'sis). *syn* sycosis. [L. *ficus*, fig]

FID Abbreviation for free induction decay.

Fiedler, Carl L.A., German physician, 1835-1921. *see* F.'s myocarditis.

field (fēld). A definite area of plane surface, considered in relation to some specific object. [A.S. *fēld*]

auditory f., the space included within the limits of hearing of a definite sound, as of a tuning fork.

Broca's f., *syn* Broca's center.

Cohnheim's f., *syn* Cohnheim's area.

f. of consciousness, *see* field of consciousness.

f. of fixation, in ophthalmology, the angular distance around which the line of fixation can be turned.

f.'s of Forel, three circumscribed, myelin-rich regions of the subthalamus known as H fields (from Haubenfelder); 1) field H₁, corresponding to the thalamic fasciculus, a horizontal fiber stratum at the junction of the subthalamus and the overlying thalamus, is composed of pallidothalamic and cerebellothalamic fibers (brachium conjunctivum) and is separated by the zona incerta from the more ventrally placed field H₂; 2) field H₂, formed by the lenticular fasciculus and arching over the dorsal border of the subthalamus, is composed largely of pallidothalamic fibers; 3) field H₃ or prerubral field, is a large field of intermingling gray and white matter immediately rostral to the red nucle-